MEDICATION CHART: Start Date:_____ End Date: _____ Symptom:_____

Name of Medicine: _____ Dose: _____

Day / Time Taken		Day / Time Taken		Day / Time Taken		Day / Time Taken	

BE INFORMED ABOUT REYE'S SYNDROME

Reye's Syndrome usually appears after a flu-like illness, upper respiratory infection, chicken pox, etc. It has been reported in children and adults.

The early signs are usually continuous vomiting, listlessness, loss of pep, aggressiveness, confusion and irrational behavior.

Medicines at the very least can mask symptoms. DO NOT give your child aspirin or anti-nausea medicine.

Phone your doctor immediately. If the doctor is unavailable, contact your closest hospital emergency team.

Abnormal Liver Test: SGOT and SGPT strongly suggest a diagnosis of Reye's Syndrome. Also there may be elevations in the blood ammonia level.

Time is important! Early diagnosis is VITAL.

(419) 636-2679 (800) 233-7393
Web: www.reyessyndrome.org Email: nrsf@reyessyndrome.org
National Reye's Syndrome Foundation, Inc.

Some Drugs Are NOT for Children!

Learn More:

www.reyessyndrome.org

reyessyndrome.wordpress.com

facebook.com/reyessyndrome

800-233-7393

National Reye's Syndrome Foundation

Other Names for Aspirin

Acetyl Salicylic Acid - ASA
Acetylsalicylate
Acetylsalicylic Acid
Aluminum Acetyl Salicylate
Ammonium Salicylate
Amyl Salicylate
Arthropan
Aspirin
Benzyl Salicylate
Butyloctyl Salicylate
Calcium Acetyl Salicylate
Choline Salicylate
Ethyl Salicylate
Lithium Salicylate
Methyl Salicylate
Methylene Disalicylic Acid
Octisalate
Octyl Salicylate
Phenyl Salicylate
Sal Ethyl Carbonate
Salicyamide
Saliclanilide
Salicylsalicylic Acid
Santalyl Salicylate
Sodium Salicylate
Stoncylate
Strontium Salicylate
Sulfosalicylic Acid
Tridecyl Salicylate
Trolamine Salicylate

Remember: Kids & Aspirin DON'T Mix!